

# AIRS/AMSU/HSB Version 6 L2 Standard Pressure Levels

Edited by:  
Edward T. Olsen



February 2013

Version 1.0



Jet Propulsion Laboratory  
California Institute of Technology  
Pasadena, CA

Submit Questions to:

<http://airs.jpl.nasa.gov/AskAirs>

# AIRS V6 L2 Standard Pressure Levels

Index	pressStd, mb	pressH2O, mb
1	1100.0	1100.0
2	1000.0	1000.0
3	925.0	925.0
4	850.0	850.0
5	700.0	700.0
6	600.0	600.0
7	500.0	500.0
8	400.0	400.0
9	300.0	300.0
10	250.0	250.0
11	200.0	200.0
12	150.0	150.0
13	100.0	100.0
14	70.0	70.0
15	50.0	50.0
16	30.0	
17	20.0	
18	15.0	
19	10.0	
20	7.0	
21	5.0	
22	3.0	
23	2.0	
24	1.5	
25	1.0	
26	0.5	
27	0.2	
28	0.1	

## AIRS V6 L2 Standard Pressure Levels

The **pressStd** array appears in the Level 2 Standard Product once per granule, and is ordered from bottom of atmosphere to top. It identifies the pressure levels upon which are reported **TAirStd**, **TAirMWOnlyStd**, **GP\_Height**, and **GP\_Height\_MWOnly**. They are also the levels that bound **O3VMRStd** and on which **O3VMRStd** and **O3VMRLevStd** are reported. The array has not been modified since launch in May, 2002.

The **pressH2O** array is also ordered from bottom of the atmosphere upward and identifies the pressure levels that bound **H2OMMRStd**, **H2OMMRSat**, and **H2OMMRSat\_liquid**. These layer quantities are reported on the pressure level that bounds the high pressure side of the layers, so that there is no 50 hPa report for these quantities. However, the level quantities, **H2OMMRLevStd**, **H2OMMRSatLevStd** and **H2OMMRSatLevStd\_liquid** are reported on all 15 pressure levels.

Both pressures arrays appear in the Level 2 Standard Product once per granule as attributes. They are shared by all retrievals, and so need not be replicated 1350 times as full swath arrays.